

REFERENCES

- [1] Agliullin M.R., Danilova I.G., Faizullin A.V., Amarantov S.V., Bubennov S.V., Prosochkina T.R., Grigor'eva N.G., Paukshtis E.A., Kuteпов B.I. (2016) Sol-gel synthesis of mesoporous aluminosilicates with a narrow pore size distribution and catalytic activity thereof in the oligomerization of dec-1-ene, *Microporous Mesoporous Mater.*, 230:118-127. <https://doi.org/10.1016/j.micromeso.2016.05.007> (in Eng.).
- [2] Wu Sh., Huang J., Wu T., Song K., Wang H., Xing L., Xu H., Xu L., Guan J., Kan Q. (2006) Synthesis, characterization, and catalytic performance of mesoporous Al-SBA-15 for tert-butylation of phenol, *Chin. J. Catal.*, 27: 9-14. [https://doi.org/10.1016/S1872-2067\(06\)60004-1](https://doi.org/10.1016/S1872-2067(06)60004-1) (in Eng.).
- [3] Martin-Aranda R.M., Cejka J. (2010) Recent advances in catalysis over mesoporous molecular sieves, *Top. Catal.*, 53:141-153.<https://doi.org/10.1007/s11244-009-9419-6>(in Eng.).
- [4] Garrone E., Fajula F. (2008) Acidity and basicity of ordered silica-based mesoporous materials, *Mol. Sieves*, 6: 213-267. https://doi.org/10.1007/3829_2007_015(in Eng.).
- [5] Birce Pekmezci Karaman, Nurbanu Cakiryilmaz, Huseyin Arbag,Nuray Oktar, Gulsen Dogu, Timur Dogu. (2017) Performance comparison of mesoporous alumina supported Cu & Ni based catalysts in acetic acid reforming. *International J. Hydrogen Energy*, 42: 26257-26269 <http://dx.doi.org/10.1016/j.ijhydene.2017.08.155> (in Eng.).
- [6] Balasubramanian V.V., Srinivasu P., Anand C., Pal R.R., Ariga K., Velmathi S., Alam Sher, Vinu A. (2008) Highly active three-dimensional cage type mesoporous aluminosilicates and their catalytic performances in the acetylation of aromatics. *Microporous and Mesoporous Mater.*, 114: 303-311. <https://doi.org/10.1016/j.micromeso.2008.01.016> (in Eng.).
- [7] Linssen T., Cool P., Vansant E.F., Aldea S., Poncelet G. (2005) Octane hydroisomerization over hexagonal mesoporous aluminosilicates synthesized from leached saponite. *J. Porous Mater.*, 12: 35-40. <https://doi.org/10.1007/s10934-005-5231-2> (in Eng.).
- [8] K-C . Park, S-K. Ihm. (2000) Comparison of Pt/zeolite catalysts for n-hexadecane hydro-isomerization, *Appl Catal*, A, 203:201-209[https://doi.org/10.1016/S0926-860X\(00\)00490-7](https://doi.org/10.1016/S0926-860X(00)00490-7)(in Eng.).
- [9] Chen C.Y., Ouyang X., Zones S.I., Banach S.A., Elomari S.A., Davis T.M., OjoA.F. (2012) Characterization of shape selective properties of zeolites via hydroisomerization of n-hexane, *Microporous and Mesoporous Mater.*, 164: 71-81. <https://doi.org/10.1016/j.micromeso.2012.07.003>(in Eng.).
- [10] Tuktin B.T., Tenizbaeva A.S., Samshat N.A., Abilmagzhanov A.Z., Shapovalov A.A. (2019) Hydroprocessing of dizeл oil fractions on modified alumina catalysts, *News of the national academy of sciences of the republic of Kazakhstan. Series chemistry and technology*, Volume 3, Number 435, 435:56-62. <https://doi.org/10.32014/2019.2518-1491.30> (in Eng.).
- [11] Locus R., Verboekend D., d'Halluin M., Dusselier M., Liao Y., Nuttens N., Jaumann T., Oswald S., Mafra L., Giebeler L., Sels B. (2018) Synthetic & Catalytic Potential of Amorphous Mesoporous Aluminosilicates Prepared by Post-Synthetic Aluminations of Silica in Aqueous Media, *ChemCatChem*, 10 (6): 1385-1397. <https://doi.org/10.1002/cctc.201701660>(in Eng.).
- [12] Antonelli D.M., Ying J.Y. (1995) Synthesis of Hexagonally Packed Mesoporous TiO₂ by a Modified Sol-Gel Method. *Angewandte Chemie International Edition in English*, 34: 2014-2017. <http://dx.doi.org/10.1002/anie.199520141>(in Eng.).
- [13] Romero A.A., Alba M.D., Klinowski J. (1998) Aluminosilicate mesoporous molecular sieve MCM-48, *J. Phys. Chem. B*, 102: 123-128. <https://doi.org/10.1021/jp971469s> (in Eng.).
- [14] Lysenko S.V., Kryukov I.O., Sarkisov O.A., Abikenova A.B., Baranova S. V., Ostroumov V.A., Kardashev S. V., Kulikov A.B., Karakhanov E.A. (2011) Mesoporous aluminosilicates as components of gas oil cracking and higher-alkane hydroisomerization catalysts. *Petroleum Chemistry*, 51: 151-156. <https://doi.org/10.1134/S0965544111030091> (in Eng.).
- [15] Karakhanov E.A., Kardashev S.V., Maksimov A.L., Baranova S.V., Kulikov A.B., Ostroumov V.A., Shirokopyas S.I., Lysenko S.V. (2012) Hydroisomerization of n-dodecane on bifunctional catalysts containing mesoporous aluminosilicates. *Petroleum Chemistry*, 52: 228-232. <https://doi.org/10.1134/S0965544112040020> (in Eng.).
- [16] Sing K.S W., Everett D.H., Haul R.A.W., Moscou L., Pierotti R.A., Rouquerol J., Siemieniewska T. (1985) Reporting Physisorption Data for Gas/Solid Systems with Special Reference to the Determination of Surface Area and Porosity, *Pure Appl. Chem.*, 57 (4): 603-619. <http://dx.doi.org/10.1351/pac198557040603>(in Eng.).
- [17] Aguado J., Escola J.M., Castro M.C., Paredes B. (2005) Metathesis of 1-hexene over rhenium oxide supported on ordered mesoporous aluminas: comparison with Re₂O₇/γ-Al₂O₃ *Appl. Catal. A: General*, 284: 47-57. <https://doi.org/10.1016/j.apcata.2005.01.017> (in Eng.).
- [18] Chakraborty B., Viswanathan B. (1999) Surface acidity of MCM-41 by in situ IR studies of pyridine adsorption, *Catal.Today*, 49: 253-260 [https://doi.org/10.1016/S0920-5861\(98\)00431-3](https://doi.org/10.1016/S0920-5861(98)00431-3) (in Eng.).
- [19] Shukla DB and Pandya VP. (1989) Estimation of crystalline phase in ZSM-5 zeolites by infrared spectroscopy. *J. Chem. Technol. Biotechnol.*, 44(2):147-154. <http://dx.doi.org/10.1002/jctb.280440206>(in Eng.).
- [20] Luis Fioravanti Isernia (2013) FTIR Study of the Relation, between Extra-framework Aluminum Species and the Adsorbed Molecular Water, and its Effect on the Acidity in ZSM-5 Steamed, *Zeolite Materials Research*, 16(4): 792-802 <http://dx.doi.org/10.1590/S1516-14392013005000044>(in Eng.).