


Kazakh Physical Society



KAZAKH  BRITISH
T E C H N I C A L
UNIVERSITY

**Second Annual Meeting
of Kazakh Physical Society**

June 6-8, 2019, Almaty

SECOND ANNUAL MEETING OF KAZAKH PHYSICAL SOCIETY

June 6-8, 2019, Kazakh-British Technical University



Chair: Prof. T. A. Kozhamkulov, President of KPS

Co-Chairs: Prof. T. S. Ramazanov, Vice-President of KPS, Al Farabi Kazakh National University,

Prof. K.A. Baigarin, Head of Nur-Sultan Branch of KPS, Nazarbayev University

Prof. M.T. Gabdullin, Head of Almaty Branch of KPS, Kazakh-British technical University

Scientific Program Committee:

KazNU: T.A. Kozhamkulov, T.S. Ramazanov, A.E. Davletov, M. E. Abishev, N.Zh. Takibayev, K.N. Dzhumagulova;

NU: K.A. Baigarin, A. Nurmukhanbetova, Z. Utegulov, A. Tikhonov, A. Desyatnikov, A. Zhumabekov, A. Bountis, M. Kaikanov;

ENU: R. Myrzakulov, A. K. Aryngazin;

KBTU: M.T. Gabdullin, K.Kh. Nussupov, N.B. Beisenkhanov, D.I. Bakranova;

KazNTU: S.E. Kumekov, R. Beisenov;

KazNPU: A.I. Kupchishin, V.N. Kosov;

NNC: E.G. Batyrbekov, M.K. Skakov;

INP: E. Kenzhin, N.T. Burtebayev, T. Zholdybayev, M. Zdorovets;

PTI: T.K. Sadykov, A. Serikkanov;

FAPI: L.M. Chechin, M. Dubovichenko;

ISR: Ch. Omarov;

II: V.M. Somsikov, O.N. Krikunova;

KSU: N. Ibrayev

TSHVNS: V.V. Zhukov

ZhSU: E.S. Andasbaev

Local Organizing Committee:

Almaty Branch of KPS

Kazakh-British Technical University



PROGRAM
SECOND ANNUAL MEETING OF KAZAKH PHYSICAL SOCIETY
Kazakh-British Technical University

Day 1
June 6, 2019

| | |
|----------------------|--|
| 08:30 - 09:30 | REGISTRATION OF PARTICIPANTS OF THE CONFERENCE LONDON ROOM (404 ROOM) |
| 09:30 - 09:50 | ANNUAL MEETING OPENING Maratbek Gabdullin , Head of Almaty Branch of KPS Tolegen Kozhamkulov , President of KPS Kanat Baigarin , Head of Astana Branch of KPS Erlan Batyrbekov , General Director of the RSE NNC RK Bum-Hoon Lee , President, Korean Physical Society |
| 09:50 - 10:10 | Bum-Hoon Lee Sogang University, Seoul, South Korea Black Holes in the dilatonic Einstein-Gauss-Bonnet theory |
| 10:10 - 10:30 | Hernando Quevedo National Autonomous University of Mexico, Mexico City, Mexico Quasi-homogeneous black hole thermodynamics |
| 10:30 - 10:50 | Tlekkabul Ramazanov Institute for Experimental and Theoretical Physics, Al-Farabi Kazakh National University, Almaty, Kazakhstan Polarization and magnetic field effects in complex plasmas |
| 10:50 - 11:10 | Sergey Maiorov Prokhorov General Physics Institute of the Russian Academy of Sciences, Moscow, Russia Joint Institute for High Temperature of the Russian Academy of Sciences, Moscow, Russia On the ion drift in cold gas |
| 11:10 - 11:30 | Anton Desyatnikov Nazarbayev University, Nur-Sultan, Kazakhstan Optical vortex rings in nonlinear media |
| 11:30 - 11:50 | Alexander Shepetov P.N.Lebedev Physical Institute (LPI), Tien-Shan Mountain Cosmic Ray Station, Moscow, Russia Simultaneous observation of lightning emission in different wave ranges of the electromagnetic spectrum in Tien Shan mountains |
| 11:50 - 12:10 | Nassurlla Burtebayev Institute of Nuclear Physics, Almaty, Kazakhstan The interaction processes of the charged particles and light ions |

| | |
|----------------------|--|
| | with the p- and sd-shell nuclei for astrophysical and thermonuclear applications |
| 12:10 - 12:30 | Alexandr Vurim NNC RK (National Nuclear Center of the Republic of Kazakhstan), Kurchatov, Kazakhstan Experimental Researches to Ensure Nuclear Power Safety at the Base of NNC RK Facilities |
| 12:30 - 14:00 | LUNCH BREAK |
| 14:00 - 14:15 | Nurzhan Beisenkhanov Kazakh-British Technical University, Almaty, Kazakhstan Physical Properties of Epitaxial Silicon Carbide Films, Grown by Atomic Substitution on the High-resistance (111) Oriented Silicon |
| 14:15 - 14:30 | Askhat Jumabekov Department of Physics, Nazarbayev University, Nur-Sultan, Kazakhstan Semitransparent Back-Contact Perovskite Solar Cells |
| 14:30 - 14:45 | Vladimir Kossov Olga Fedorenko Abai Kazakh National Pedagogical University, Almaty, Kazakhstan Special Regimes under Diffusion in Gaseous Mixtures |
| 14:45 - 15:00 | Askar Davletov Al-Farabi Kazakh National University, Almaty, Kazakhstan Roton structure of dust-acoustic wave spectrum of complex plasmas with finite-size particles |
| 15:00 - 15:15 | Vladimir Dzhunushaliev Institute for Experimental and Theoretical Physics, Al-Farabi Kazakh National University, Almaty, Kazakhstan Non-Abelian Proca-Dirac-Higgs theory: particle-like solutions and a mass gap |
| 15:15 - 15:30 | Vladimir Messerle Combustion Problems Institute of Ministry of Education and Science of Kazakhstan, Almaty, Kazakhstan Plasma Application for Uranium-Containing Solid Fuels Processing |
| 15:30 - 16:00 | COFFEE BREAK |
| 16:00 - 16:15 | Nurgali Takibayev al-Farabi Kazakh National University, Almaty, Kazakhstan Induced Processes and Reactions in Neutron Star Crusts |
| 16:15- 16:30 | Alma Dauletbekova L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan Synthesis of nanocrystals in SiO₂/Si track template by |

Special Regimes under Diffusion in Gaseous Mixtures

¹Kossov V.N., ²Fedorenko O.V., ²Mukamedenkyzy V.

¹Abai Kazakh National Pedagogical University, 13 Dostyk Ave., Almaty, Republic of Kazakhstan

²Al-Farabi Kazakh National University, 71 al-Farabi Ave., Almaty, Republic of Kazakhstan

e-mail: kosov_vlad_nik@list.ru

Abstract: The features of the occurrence of convective flows during diffusion in multicomponent gas mixtures are considered. Experimental data and results of a numerical analysis of the emergence of convective motions during diffusion in three-component gas mixtures are presented. It is shown that the analysis of the stability of three-component gas mixtures can be carried out both in the framework of the linear stability theory and modern methods of numerical simulation.

1. Introduction

Molecular diffusion can lead to instability of the mechanical equilibrium of the mixture with the subsequent occurrence of natural convection, which noticeably intensifies the total mass transfer. In experiments on the study of mixing in multicomponent systems at different pressures and compositions [1], and diffusion of a mixture of solution vapors into an inert gas [2], convective flows were observed leading to a synergistic effect associated with a significant increase in the rate of mixing of the system components.

The solution of issues related to the determination of the mechanism of change of the regimes "diffusion – convection" in multicomponent gas mixtures and the parameters determining the occurrence of the convective mode are important for fundamental and applied problems of convective mass transfer.

2. Experimental data

In the ternary gas mixtures (for example $0.5538 \text{ H}_2 + 0.4462 \text{ N}_2 - \text{CH}_4$ and $0.8846 \text{ H}_2 + 0.1154 \text{ CH}_4 - \text{He}$ [3], $0.5504 \text{ CH}_4 (1) + 0.4496 \text{ Ar} (2) - \text{N}_2 (3)$, $0.5143 \text{ He} (1) + 0.4857 \text{ Ar} (2) - 0.5148 \text{ CH}_4 (3)$ and $0.8366 \text{ CH}_4 (1) + 0.1634 \text{ R12} (2) - n\text{-C}_4\text{H}_{10} (3)$ [4]) a regime change can occur, due to the difference in diffusion coefficients. The parameters defining such a transition are the pressure and the initial composition of the mixture [3, 4]. In this case, it is possible to realize the conditions when the transfer of the component with the highest molecular weight will be prior to the others. This situation that is not typical for the diffusion can be explained by the occurrence of convective structured formations.

3. Numerical analysis

Diffusion mixing near the boundary of the change of kinetic regimes can be described by the general system of hydrodynamic equations, which includes the Navier-Stokes equations written in the Boussinesq approximation, preserving the number of particles of a mixture and components [1, 3, 4]. This system of equations can be solved analytically using the linear stability theory [3] or by means of the splitting scheme for physical parameters [4].

First approach allows in terms of diffusion Rayleigh numbers to obtain the maps of stability for the ternary gas mixtures determining regime change "diffusion – concentration gravitational convection" for the vertical channels. Thus, in order to evaluate the theoretical effect of pressure (or other influencing parameter) on the occurrence of convection under the isothermal mixing of ternary gas mixture is necessary to consider the location of the partial Rayleigh numbers (as points) relative to the boundary line on the plane $(\text{Ra}_1, \text{Ra}_2)$. If the partial Rayleigh numbers lie below the boundary line, then the diffusion process is observed in the system. If the partial Rayleigh numbers are in the area between the boundary line and zero density gradient line, then the stability paradox occurs in the system, i.e. there is an unstable diffusion mixing. If the partial Rayleigh numbers are situated above the boundary line, then the convective mixing is observed in the mixture.

The numerical model for the study of isothermal transfer in three-component gas mixtures permits to determine and analyse the behaviour of isoconcentration lines of the components. The main feature of the transition from the diffusion mode to the convective mode is the appearance of nonlinear isoconcentration lines due to a significant difference in the coefficients of mutual diffusion of the components. With increasing pressure or content in the mixture of the component with the highest molecular weight, the curvature of the isoconcentration lines increases which is the cause of the occurrence of convective instability. With unstable mixing, a pulsating transfer mode is possible, which is associated with the occurrence of structural convective formations.

4. Summary

Thus, the proposed mathematical models for the study of isothermal transfer in three-component gas mixtures allows for determining the conditions under which the change of regimes "diffusion – concentration gravitational convection" occurs.

[1] V. N. Kossov, D. U. Kul'zhanov, Y. I. Zhavrin, O. V. Fedorenko, "Emergence of convective flows during diffusional mass transfer in ternary gas systems: The effect of component concentration", *Russ. J. Phys. Chem. A* 91 (6), 984 (2017)

[2] V. V. Dil'man, V. A. Lotkhov, "Molecular turbulent evaporation in a gravitational field", *Theor. Found. Chem. Eng.* 49 (1), 102 (2015).

[3] V. Kossov, S. Krasikov, O. Fedorenko, "Diffusion and convective instability in multicomponent gas mixtures at different pressures", *Eur. Phys. J. Spec. Top.* 226 (6), 1177 (2017).

[4] V. Kossov, O. Fedorenko, D. Zhakebayev, "Features of multicomponent mass transfer in gas mixtures containing hydrocarbon components", *Chem. Eng. Technol.* 42 (4), 896 (2019).