

ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В.СОКОЛЬСКИЙ АТЫНДАҒЫ «ЖАНАРМАЙ,
КАТАЛИЗ ЖӘНЕ ЭЛЕКТРОХИМИЯ ИНСТИТУТЫ» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

АО «ИНСТИТУТ ТОПЛИВА, КАТАЛИЗА И
ЭЛЕКТРОХИМИИ ИМ. Д.В. СОКОЛЬСКОГО»

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

JSC «D.V. SOKOLSKY INSTITUTE OF FUEL,
CATALYSIS AND ELECTROCHEMISTRY»

ХИМИЯ ЖӘНЕ ТЕХНОЛОГИЯ СЕРИЯСЫ



СЕРИЯ ХИМИИ И ТЕХНОЛОГИИ



SERIES CHEMISTRY AND TECHNOLOGY

5 (431)

**ҚЫРКҮЙЕК – ҚАЗАН 2018 ж.
СЕНТЯБРЬ – ОКТЯБРЬ 2018 г.
SEPTEMBER – OCTOBER 2018**

1947 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1947 ГОДА
PUBLISHED SINCE JANUARY 1947

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р ы
х.ғ.д., проф., ҚР ҰҒА академигі **М.Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

Ағабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бүркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф. корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҰҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / chemistry-technology.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Редакционная коллегия:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф. чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф. академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Гаджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz / chemistry-technology.kz>

© Национальная академия наук Республики Казахстан, 2018

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,
Институт органического катализа и электрохимии им. Д. В. Сокольского,
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail: orgcat@nursat.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov**

E d i t o r i a l b o a r d :

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadjikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.
ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/chemistry-technology.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Editorial address: Institute of Organic Catalysis and Electrochemistry named after D. V. Sokolsky
142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22,
e-mail: orgcat@nursat.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2018.2518-1491.10>

Volume 5, Number 431 (2018), 74 – 81

A.Zh. Kaldybekova, A.T. Amangazyeva, Z.B. Halmenova, A.K. Umbetova

Kazakh National University named after al-Farabi, Almaty, Kazakhstan
A_Arai_95@mail.ru, zaure.halmenova@mail.ru, alma_0875@mail.ru, amangazyeva.a@mail.ru

**DEVELOPMENT OF TECHNOLOGY FOR
THE COMPLEX ISOLATION OF BIOLOGICAL ACTIVE SUBSTANCES
FROM PLANTS OF THE GENUS HAPLOPHYLLUM A. JUSS**

Abstract: In this study, a complex study of plants of the genus *Haplophyllum* A.Juss of a sort *H. Perforatum* Kar et Kir. The main parameters or raw material quality: preparation raw material (grinding, sieve analysis); humidity, total ash, ash is insoluble 10 % HCl, sulphate ash, extractives.

Macro- and microelement composition of total ash is analyzed. Qualitative reactions and chromatographic analysis methods, the main classes of natural compounds in raw materials and their quantitative contents have been identified.

Keywords: Rutaceae, *Haplophyllum*, moisture, ash, coumarins, *Perforatum*, flavonoids, terpenoids.

Introduction

The stirpes Rutaceae includes 150 genera and about 900 species are widely distributed in tropical, subtropical and heat-sensitized regions of both hemispheres. Most of the representatives of the stirpes grow in the South of the USA (California, Florida) in the Mediterranean countries (Spain, Italy, Morocco, etc.), Brazil, Argentina, Japan, China, South Africa and the arid regions of Australia [1; 201].

Rutaceae - occur mainly as evergreen trees or bushes, sometimes lianas, very rarely perennial (*ruta - Ruta*, ash - *Dictamnus*, *haplophyllum - Haplophyllum*) and annual grasses [2; 240].

Plants of the genus *Haplophyllum* A. Juss (stirpes Rutaceae) on a global scale are represented by approximately 70 species, which are distributed from the Mediterranean to Western Siberia. There are 23 species of *Haplophyllum* growing on the territory of the Central Asian countries [3; 229].

The main active components of the plant genus *Haplophyllum* A. Juss are:

- coumarins (*collinin*, *obtusifol*, *obtusinin*, *obtusinol*, *obtosozid*, *daurosod* D, *ostol*, 6-methoxy-7-(3',3'-dimethylallyloxy) coumarin, 5-hydroxy-7-methoxycoumarin, *scopoletin*, *isoskopoltin*, *bungenediol*, *skimianin*, *umbelliferone*, and *daurozides* A and B);

- flavonoids (*haplozid* A, 7-O-glycoside *haplogenin*, *haplozid* D, *haplozid* F, 7-O-β-D- (6acetyl-), *glucopyranoside* 3,5,4'-trihydroxy-8,3' - dimethoxyflavone, 7-O-α-L-rhamnopyranosyl, glycosides of *quercetin*, *kaempferol*, *isoramnetin*, 7-glucoside of 3'-methyl ester of *gossipetin*);

- alkaloids (*acetamide*, *folifin*, *evoxin*, *haplotin*, *candecin*, *palmatin*, *magnophlarin*, *haplophin*, *buharaine*, *bucharidine*, *robustin*, *folimin*, 4-methoxy-N-methylquinolone-2, *robustinine*, *dictamnin*, *skimianin*, *dubinin*, *dubamine*, *haplopherin*, *foliosin*);

- terpenoids;

- steroids (*sitosterol*, *stigmasterol*, *camperterol*);

- essential oils;

- carboxylic acids;

- lignans (*eudesmine*, *variantcoside*, *diphyllin*, *pluviatil*, *suicilactone*, *yusticidin* B, *daurinol*);

- tannins;

- amides;

- carbohydrates (*glucose*, *galactose*, *mannose*, *xylose*, *arabinose*, *rhamnose*);

- saponins;
- higher aliphatic alcohols (n-triacontanol) and a number of other BAS (Biologically active substances) [4; 13].

Table 1 - the biologically active substances of plants of the genus *Haplophyllum* [5; 6].

Kind of plant	Dedicated connections		
	Coumarins	Flavonoids	Lignans
<i>Haplophyllum Perforatum Kar et Kir</i>	scopoletin, scopiolin, haploperoside A, haploperoside B, x haploperoside C, haploperoside D, haploperoside E	haplogenin, limocitrine, limocitrine-7-O- β -D- (6 "-O-acetyl) -glucopyranoside, haplozid A, haplozid B, haplozid C, haplozid D, haplozid E	eudesmine
<i>Haplophyllum Obtusifolium Ledeb</i>	scopoletin, 6-methoxy-7- γ , γ -dimethylal-liloxicoumarin, obtusinol, obtusin, otosozid, feruloylskopolin fraxetin, kapensin, obtusicin, obtusiprenin, otusiprenol, obtusidine, obtusipholine, fraxetine-7-O-glucoside		yusticidin B, diphyllin
<i>Haplophyllum Davuricum (L.) G. Don</i>	umbelliferon, scopoletin, 5,7-dihydroxyquarium, skimmin, daurozide A, daurozide B, daurozide D	haplogenin, haplozid B, haplozid D	yusticidin B, diphyllin daurinol
<i>Haplophyllum Latifolium Kar et Kir</i>	-	haplogenin, haplozid B, haplozid D	-
<i>Haplophyllum Versicolor Fisch. Et Mey</i>	-	-	variantcoside

From the data presented in Table 1, the common feature of all the species of studied for the content of coumarins is the presence of scopoletin, scopolin, haploperoside *Haplophyllum Perforatum Kar et Kir*, haploperoside B, haploperoside C, haploperoside D, haploperoside E. The content of flavonoids: haplogenin, limocitrine, limocitrine-7-O- β -D- (6 "-O-acetyl) -glucopyranoside, haplozid A, haplozid B, haplozid C, haplozid D, haplozid E. Content of lignans: eudemin.

Table 2 - the quantitative content of coumarins during flowering and fruiting [5; 7].

Coumarins	The yield of coumarins (in% of the weight of an air-dry plant)	
	Flowering period	Fruiting period
scopoletin,	-	trace amounts (<0,004%)
6-methoxy-7- γ , γ -dimethyl-lyloxy coumarin	0,023	trace amounts (<0,004%)
Obtusinol	trace amounts (<0,004%)	-
Obtusoside	0,08	0,19
Feruloylskopolin	0,054	0,27
Thraxetine	-	0,005
Capensin	0,018	0,14
Obtusicin	0,84	0,17
Obtusiprenin	0,025	0,26
Obtusiprenol	-	0,023
Obtusidine	-	0,025
Thraxetine-7-O-glucoside	-	0,013
	-	0,006

The data obtained by us and other researchers show that the qualitative and quantitative compositions of coumarins of the studied plant species vary greatly depending on the ecological-geographic, soil-

climatic conditions, as well as the vegetation period and the plant organ. The same species of this genus, growing in different geographical areas, may contain different in quality and quantity composition of coumarins.

As can be seen from the data presented in Table 2, the greatest amount of coumarins is extracted during the flowering period, rather than during the fruiting period, since during the fruiting period the yield of coumarins decreases.

In the flora of Kazakhstan, there are more than 70 species of plants, many species have plant medicinal plants and may be required for the pharmaceutical industry of the Republic. Therefore, the study of the chemical composition of pensions is not only scientific, but also practical [4; 12].

Various species of *Haplophyllum* (*H. Dubium*, *H. Bucharicum*, *H. Bungei*, *H. Davuricum*, *H. Foliosum*, *H. Obtusifolium*, *H. Latifolium*, *H. Perforatum* and others) have long been used in folk medicine for the treatment of skin, as well as an antidote for poisoning, antipyretic, analgesic, laxative, for diseases of the stomach and spleen. Extracts of some species show antitumor and cytotoxic activity. Therefore, they attract the attention of researchers as potential sources of biologically active substances. The objects of our study are the species *Haplophyllum Perforatum* [4; 14].

In plant resources under the editorship of Sokolov P.D. information on the chemical composition and useful properties of this plant species is presented. Phenyl carboxylic acids and their derivatives: methyl ester of n-coumaric (n-hydroxycinnamic) acid. Lignans: (8R, 8'R 4, 4'-dimethylglylignolide-9.9 ". The plant roots contain alkaloids (in %) 0.047, skimianin 0.009, evoxin, haplopin. Containment in the aerial part of plants: steroids (in %) sitosterol 0.07; alkaloids (in %) 0.04-0.05; skimianin 0.14, haplatin 0.001, evoxin 0.04, haplopin 0.001, haplamide 0.003, haplamidine, glycopein, 7-isopentenyl-oxy- γ -pharagin. Phenyl carboxylic acids and their derivatives: methyl ester of n-hydroxycinnamic (n-coumaric) acid. In the stems of the plant contain: alkaloids 0.23%. In the leaves of the plant contain: alkaloids (in %) 0.04-0.6, skimianin 0.005-0.6, hapsolin 0.06, dubamine, haplatin.

Useful properties. The content of n-hydroxycinnamic acid methyl ester in the aerial part of plants shows estrogenic activity [5; 13].

And in plant resources under the editorship of Budantsev A.L. information on the chemical composition and useful properties of this plant species is also presented. The roots contain:

- terpenoids (fraksinellon);
- alkaloids (robustin, dictamnin, γ -pharagin, skimianin, evoxin, haplozidine 0.01%, haplosinin 0.003%);
- coumarins (seselin);
- flavonoids (haplogenin, limocitrin, haplozid C).

The leaves of the plant contain:

- flavonoids (7,3'-dimethyl ether of dihydroquercetin, 3,6-dimethyl ester of quercetin (axillarin), 3,5,7-trimethyl ester of kaempferol, 7-O- β -D-methyl glucuronide of 3'4'-dimethyl ester of lutheline, 7,3'-dimethyl ether and 3'-methyl esters of luteolin).

In the flowers and fruits are found essential oil in its composition includes monoterpenoids, hydrocarbons.

Water extract is used externally for dermatoses, eczema, varicose veins, alopecia, asthenia, anemia [6; 234,235].

Result and Discussion

The size of the medicinal plant material is determined by the method of sieve analysis.

The investigated plants were crushed to a particle size within 0.5mm in flowers (420.759), 1mm in stems (316.488), according to the regulatory documentation.

Good quality - compliance of medicinal raw materials, products, medicines with technical requirements of NTD.

All indicators of good-quality were determined by the methods of the GF RK, the European Pharmacopoeia and other literary sources.

Table 3 - Sieve analysis of H.Perforatum plant

Sifter №	Mass (<i>H.Perforatum</i>) = 600g		Mass (<i>H.Perforatum</i>) = 500g	
	<i>H.Perforatum</i> (flowers)		<i>H.Perforatum</i> (stems)	
	Mass, g	%	Mass, g	%
10	-	-	-	-
8	-	-	-	-
6	9,545	1,59	21,352	4,27
4	19,696	3,28	27,640	5,53
2	35,348	5,89	40,552	8,11
1	72,443	12,07	316,488	63,3
0,5	420,759	70,13	28,263	5,65
0,3	12,928	2,15	24,499	4,9
0,2	9,863	1,64	16,020	3,2
0,25	6,454	1,08	11,125	2,23
0,1	5,132	0,86	7,085	1,42
Remain	3,552	0,59	2,120	0,42
Losses	4,280	0,71	4,856	0,97

Table 4 - Numerical indicators of benignity of a plant of H. Perforatum species

Indicators of good quality	Plant H. Perforatum	
	Flowers	Stems
Humidity, %	5,74	5,83
Total ash, %	6,74	6,38
Sulphated ash, %	11,25	12,43
Ashes insoluble in 10% HCl	9,56	10,18
Containment of detective substances (70% alcohol)	44,06	32,59

As can be seen from the data presented in table 3, the moisture content in the stems (5.83%) is greater than for the flowers of the plant (5.74%). The total ash of the plant was 6.74%. The ash insoluble in 10% HCl was 9.56% in flowers, 10.18% in stems. And the distribution of extractive substances in the plant was H. Perforatum in flowers 44.06%, stems 32.59%.

The solvent, which should be taken when extractives are determined, is indicated in the relevant specification for this type of raw material. Usually, it is the same solvent that is used when preparing a tincture or extract from this raw material.

Extractive substances show the content of the likely active substances in the raw materials, which can later be processed. As can be seen from the data, the greatest amount of extractive substances is extracted by the flowers of this plant. The obtained data testify to the quality of the raw materials and the profitability of using the BAS complex. Figure 1 shows a benign chart of the distribution of H. Perforatum Kar.et Kir.




Figure 1 - Diagram of the assimilation of Haplophyllum Perforatum.

In the plant raw material, the ash of total, sulfate ash, ashes is insoluble in 10% HCl, which is the residue after treatment with the total HCl ash and consists mainly of silicates, which are a natural component for some objects, but more often the result of contamination of the raw materials with sand, earth and stones.

Mineral elements by their content in the plant are divided into macro elements, microelements and ultra microelements. Macroelements include Na, K, Ca, Mg, Cl, P; their content in the ashes is measured in hundredths of a percent. Microelements: Zn, Cu, Ni, Mn, Cu. Along with the "biometals" - elements that make up living organisms (for example, sodium, potassium, calcium, magnesium, iron, zinc), they can be referred to heavy metals - cadmium, lead, chrome, mercury and other d-elements, content which, according to the gradation of Vinogradova A.P., corresponds to the level of microelements in plants.

The study was carried out by the method of atomic absorption analysis on the basis of the Center for Physical and Chemical Methods of Research and Analysis (CPCMRA). The data are presented in table 5.

Table 5 - Macro- (K, Ca, Na, and Mg) content and microelements (Cu, Cd, Pb, Fe, Mn, Ni, Zn) in the plant *H. Perforatum*

Microelements	Flowers of a plant, %	The stems of the plant, %
Cd	$0,0227 \cdot 10^{-3}$	0
Pb	$0,1636 \cdot 10^{-3}$	$0,1487 \cdot 10^{-3}$
Fe	$5,2048 \cdot 10^{-3}$	$3,3864 \cdot 10^{-3}$
Mn	$0,6036 \cdot 10^{-3}$	$0,2767 \cdot 10^{-3}$
Ni	$0,2569 \cdot 10^{-3}$	$0,2301 \cdot 10^{-3}$
Zn	$2,7284 \cdot 10^{-3}$	$208,680 \cdot 10^{-3}$
Cu	$0,3671 \cdot 10^{-3}$	$0,3691 \cdot 10^{-3}$
Macroelements		
K	$473.370 \cdot 10^{-3}$	$432.30 \cdot 10^{-3}$
Ca	$395.880 \cdot 10^{-3}$	$157.890 \cdot 10^{-3}$
Na	$23.5525 \cdot 10^{-3}$	$28.2150 \cdot 10^{-3}$
Mg	$91.180 \cdot 10^{-3}$	$45.7025 \cdot 10^{-3}$

The largest number of macro and microelements is the flowers of the plant. In the stems and flowers of the plant, the dominant macro elements are K. In the flowers and stems of the plant, there is an increased content of macroelements like K, Ca. And the content of macroelements K, Ca, Na is explained by the fact that in saline conditions the Na +, K + concentration is increased, and they diffuse rapidly into the interior and easily saturate the cell sap.

A comparative phytochemical analysis for qualitative reactions and chromatographic analysis for the main classes of biologically active substances was carried out. The results are shown in Table 6.

Table 6 - Phytochemical analysis of the plant *Haplophyllum A. Juss*

BAS	Developers	Flowers	Stems
Carbohydrates	o-Toluidine	Light-brown envelope	Light-brown envelope
Tannins	JAK	Blue shading	Black-blue staining
Phenolic compounds	FeCl ₃ , 3%	Green	Dark green
Flavonoids	NH ₃ , 5%	bright yellow	yellow-green
	AlCl ₃	pale yellow	bright yellow
Carotenoids	SiHNO	-	-
	AgNO ₃ , 1%	reddish-brown	reddish-brown
	NaOH, 5%	-	-
Amino acids	KMnO ₄ (Potassium permanganate)	decolorization	decolorization
	Phosphorus-molybdcic acid	blue	green
Carboxylic acids	C ₉ H ₆ O ₄	pale purple	light purple
	(ninhydrin)	yellow-brown	brown
	CH ₄ N ₂ O (urea)	-	-
	MgAc ₂	-	-

As can be seen from the data presented in table 6, in the accumulation of *Haplophyllum A. Juss* there are the main groups of biologically active substances. Carbohydrates, phenolic compounds, tannins, flavonoids, carotenoids, amino acids and carboxylic acids were found.

The method of the two-story paper chromatography in the use of reliable samples in flowers, and packages from flavonoids were occupied by quercetin and rutin; from the corners – cane sugar and glucose; from amino acids - as the so-called and all-alone α -amino acids.

An important parameter for obtaining the plant extract is the ratio of raw materials and solvent from 1:4 to 1:8. For 5 grams of raw material (flowers and stems of *H. Perforatum Kar.et Kir*) Pour in different volumes and different percentages of ethyl alcohol, when using a curing agent 1:6, and instill 24 hours without decontamination at a temperature of 24-25°C. These parameters are presented in table 7.

Table 7 - Determination of the optimum extragent for extraction

Mass of raw material, g <i>const</i>		5	5	5
<i>Solvents (extragents)</i>		30% ethanol	50% ethanol	70% ethanol
<i>Extraction temperature, °C const</i>		24-25°C	24-25°C	24-25°C
Weight of dry extract, g	Flowers	0,1253	0,1625	0,0941
	Stems	0,3442	0,5661	0,3824
Extraction amount, %	Flowers	2,51	3,25	1,88
	Stems	6,88	11,32	7,65

As a result of comparative studies it was found that the optimal extragent is 50 % ethyl alcohol for flowers, 70 % ethyl alcohol for stems. When the ratio of raw materials: extragent 1: 6, the amount of dry extract in flowers was 3.25 %, and in stems 7.65 %.

One of the important parameters in the technology of obtaining extracts is also the ratio of the selected extragent to the raw material. To determine the optimum volume, the ratio of raw materials and solvent is changed from 1:4 to 1:8, pouring on 5 g of vegetable raw materials 70 % and 50 % ethyl alcohol. At this time constant of extraction (24 hours) and temperature (24-25° C), the following results were obtained. This is presented in the form of a table 8.

Table 8 - Determination of the optimal ratio

Mass of raw material, g <i>const</i>		5	5	5
Ratio of raw materials (g) and extraction (ml)		1:4	1:6	1:8
<i>Extraction temperature, °C const</i>		24-25°C	24-25°C	24-25°C
Weight of dry extract, g	Flowers	0,291	0,1625	0,2462
	Stems	0,1001	1,234	0,3824
Extraction amount, %	Flowers	5,82	3,25	4,92
	Stems	2,002	24,68	7,65

The data presented on the table show that the ratio of 1:6 with the extragent chosen was optimal, in flowers the amount of extract was 3.25%, and in stems 7.65%.

Table 9 - Determination of the optimal time

Mass of raw material, g <i>const</i>		5	5	5
Extraction time		12	24	48
<i>Extraction temperature, °C const</i>		24-25°C	24-25°C	24-25°C
Weight of dry extract, g	Flowers	1,618	0,1625	1,1125
	Stems	0,9864	1,8515	0,3824
Extraction amount, %	Flowers	32,36	3,25	22,25
	Stems	19,73	37,03	7,65

As a result, it was found that the optimal time was 24 hours with the extragent selected, the mass of the dry extract was 0.11625g in flowers and 0.3824g in stems.

The carried out researches allowed to determine the optimum values of the parameters influencing the extraction of biologically active substances from the plant *H. Perforatum* Kar.et Kir. stirpes Rutaceae.

Conclusions

Based on the results of the study, the following conclusions can be drawn:

The numerical parameters of the raw materials are determined: sieve analysis, moisture content, total ash, fly ash soluble in 10 % HCl, sulfate ash, extractives regulating the quality of raw materials complies with the norms of the State Pharmacopoeia of the Republic of Kazakhstan.

Methods of spectral analysis established for the plant *H. Perforatum* 4 macro- and 7 microelements. Qualitative methods of analysis include carbohydrates, tannins, flavonoids, phenolic compounds, carotenoids, amino acids and carboxylic acids.

The technology of extraction of the extract from the studied type of raw material, its ratio, temperature and time is worked out. The following parameters were chosen as the optimal condition: extragent 50 % ethyl alcohol in flowers, 70 % ethyl alcohol in stems, extraction ratio 1:6, extraction time 24 hours, temperature 24-25°C.

REFERENCES

- [1] Vvedenskij i A.I. Sem. LXXXIII. Rutovy'e – Rutaceae. // Flora SSSR: v 30 t./ nachato pri ruk. i pod. gl. red. B. L. Komarova. – M. – L.: Izd-vo AH SSSR, 1949. T. XIV/red. toma B.K. SHishkin, E.G. Bobrov. S. 198. 24b. 790s. 4000 e'kz.
- [2] Rodionenko G.I. Sem. 38. Rutovy'e – Rutaceae. //Derev'ya I custarniki SSSR. Dikorastushhie, kultiviruemy'e I perspektivny'e dlya introdukczii./Red. toma. S.YA.Sokolov. M. L.:Izd-vo AH SSSR, 1958 – T.IV. Pokry'tosemny'e. Semejstva Bobovy'e –Granatovy'e. S.230 – 256. 976s. 2500e'kz.
- [3] Vvedenskij A.I. Rod845. Yasenez – Dictamnus L./Flora SSSR: v 30 t./nachato pri ruk. i pod gl. red. V.L.Komarova. M. L.: Izd-vo AH SSSR, 1949. T.XIV/red. toma B.K. SHishkin, E.G. Bobrov. S. 227 – 232. 790 s. 4000e'kz.
- [4] E'.X. Botirov, M.P. Yuldashev, A.D. Matkarimov, V.M. Malikov, //Kumariny', flavanoidy' i lignany' pyati vidov rastenij roda Haplophyllum A. Juss. //Ximiya rastitel'nogo sy'rya. 2015. №1. S. 5-14.
- [5] П.Д. Соколов Растительные ресурсы СССР/Цветковые растения, их химический состав, использование. Академия наук СССР Ботанический Институт им.В.Л. Комарова, Санкт-Петербург. 1991. 58-60 с.
- [6] A.L. Budanczev. Sem. Lycopodiaceae-Ephedraceae.//rastitel'nogo resursy//Chast' I. Rossijskaya Akademiya Nauk Botanicheskij Institut im. V.L. Komarova, Sankt. Peterburg.1996.S.234-235.

**А.Ж. Қалдыбекова, А.Т. Амангазиева,
З.Б. Халменова, А.К. Үмбетова**

Әл-Фараби атындағы Қазақ Ұлттық Университеті, Алматы, Қазақстан

НАРЛОПHYLLUM A. JUSS ШӨБІНЕН БИОЛОГИЯЛЫҚ БЕЛСЕНДІ ЗАТТАРДЫҢ КЕШЕНДІ БӨЛІНУ ТЕХНОЛОГИЯСЫН ДАМЫТУ

Аннотация. Бұл зерттеуде Rutaceae тұқымдасына жататын *Haplophyllum A.Juss* өсімдігінің *H.Perforatum Kar et Kir* түріне комплексті зерттеу жүргізілді. Шикізат сапалылығының маңызды параметрлері қаралды. Олар: шикізат дайындау (майдалау, електен өткізу зерттеуі); ылғалдылық, жалпы күлділік, 10 % HCl-да ерімейтін күлділік, сульфатты күлділік, зерттелініп отырған өсімдік түрлерінен экстрагенттің табиғатын, оның шикізатпен қатынасын, экстрактілеу уақыты мен жиілігін өзгерте отырып, биологиялық белсенді кешенді алу технологиясы өңделді.

Жалпы күлділіктің макро- және микроэлементтік құрамы зерттелді. Сапалық реакция және хроматографиялық талдау әдісімен шикізаттағы табиғат қосылыстарының маңызды класстары жіктелді және олардың сапалық құрамы анықтады.

Түйін сөздер: Rutaceae, Haplophyllum, ылғалдылық, күлділік, кумариндер, Perforatum, флаваноидтар, терпеноидтар.

А.Ж. Калдыбекова, А.Т.Амангазиева, З.Б. Халменова, А.К. Умбетова

Казахского Национального Университета имени аль-Фараби, Алматы, Казакстан

РАЗРАБОТКА ТЕХНОЛОГИИ КОМПЛЕКСНОГО ВЫДЕЛЕНИЯ БИОЛОГИЧЕСКИХ АКТИВНЫХ ВЕЩЕСТВ ИЗ РАСТЕНИЙ РОДА *HAPLOPHYLLUM* A. JUSS

Аннотация. В данном исследовании проведено комплексное исследование растения рода *Haplophyllum* A.Juss вида *H. Perforatum* Kar et Kir. Рассмотрены основные параметры качества сырья: подготовка сырья (измельчение, ситовой анализ); влажность, общая зола, зола не растворимая в 10 % HCl, сульфатная зола, экстрактивные вещества.

Проанализирован макро- и микроэлементный состав общей золы. Качественными реакциями и хроматографическими методами анализа идентифицированы основные классы природных соединений в сырье и определено их количественные содержание. В сырье определены основные технологические параметры получения экстракта.

Ключевые слова: Rutaceae, *Haplophyllum*, влажность, зольность, кумарины, *Perforatum*, флаваноиды, терпеноиды.

Information about authors:

Kaldybekova Araylym Zhunibekyzy - Second year master of the Kazakh National University named after al-Farabi faculty of "Chemistry and Chemical Technology";

Khalmenova Zaure Beisentaevna - Ph.D. Associate Professor of Organic Chemistry and Technology of Organic Substances of the Chemistry of Natural Compounds and Polymers;

Umbetova Almagul Kendebaevna - Ph.D. Associate Professor of Organic Chemistry and Technology of Organic Substances of the Chemistry of Natural Compounds and Polymers;

Amangazieva Asem Talgatovna - A fourthyear student of the Kazakh National University named after al-Farabi faculty of "Chemistry and Chemical Technology".

МАЗМҰНЫ

<i>Кантуреева Г.О., Defrancesco E., Алибеков Р.С., Уразбаева К.А., Ефимова И.Е.</i> Қазақстанның дәстүрлі азық-түлік өнімдерді сәйкестендіру жаңа тенденциялары	6
<i>Туктин Б.Т., Теңізбаева А.С., Нұрғалиев Н.Н., Шаповалова Л.Б., Яскевич В.И.</i> Модифицирленген Ni(Co)-Mo- Al ₂ O ₃ катализаторларында тура айдалған бензин фракциясын гидроизомерлеу және гидроөңдеу	13
<i>Ахметалимова А.М., Ивасенко С.А., Марченко А.Б., Ишмуратова М.Ю., Полезчак Э., Людвичук А., Посева И.В.</i> Қарағанды өңіріндегі <i>THYMUS EREMITA</i> KLOK. және <i>THYMUS RASITATUS</i> KLOK. өсімдіктерінің химиялық құрамын зерттеу.....	20
<i>Фазылов С.Д., Нұркенов О.А., Журинов М.Ж., Әрінова А.Е., Туктаров А.Р., Исәева А.Ж., Шаихова Б.К.</i> С ₆₀ фуллеренге гидразондардың палладий комплекстерімен катализденетін циклооксидің синтезі	26
<i>Опимах Е.В., Левданский А.Э., Голубев В.Г., Корганбаев Б.Н., Сарсенбекулы Д.</i> Ұсақтау барысындағы меншікті энергия шығындарын төмендетудің келешекті бағыттары	32
<i>Қапсәмет М.Ж., Тәжібаева С.М., Уракаев Ф.Х., Уралбеков Б.М., Бүркімбаев М.М., Бачилова Н.В.</i> Нанокүкіртті алу және тұрақтандыру	41
<i>Байсанов С.О., Толоконникова В.В., Нарикбаева Г.И., Корсукова И.Я., Жучков В.И.</i> Күй диаграммасына талдау жасау негізінде марганецті және хромды феррокорытпаларды балқытуға термодинамикалық бағалау.....	47
<i>Құлқасымов Ж.Ә., Нұртаева Г.Қ., Мұстафин Е.С., Айнабаев А.А., Мұстафин Т.Е., Борсынбаев А.С., Жарикесов Ф.А.</i> Теңізге төгілген мұнайды жоюда хердерлерді пайдаланудың мүмкіндіктері	58
<i>Туктин Б.Т., Нұрғалиев Н.Н., Тенизбаева А.С., Шаповалова Л.Б., Комашко Л.В.</i> Бензиннің әртүрлі фракцияларын модифицирленген алюмокобальтмолибден катализаторларында гидрожақсарту	67
<i>Қалдыбекова А.Ж., Амангазиева А.Т., Халменова З.Б., Үмбетова А.К.</i> <i>Harporhyllum</i> A. Juss шөбінен биологиялық белсенді заттардың кешенді бөліну технологиясын дамыту	74
<i>Опимах Е.В., Левданский А.Э., Волненко А.А., Жумадуллаев Д.К.</i> Флотациялық процесстерді жүргізу әдістері	82
<i>Чиркун Д. И., Левданский А. Э., Волненко А.А., Сарсенбекулы Д.</i> Соккылы-ортдан тепкіш диірмендердегі бөлшектердің динамикасын зерттеу	92
<i>Баймұқашева Г.К., Қалауова А.С., Құспанова Б.К., Насиров Р.Н.</i> Үшфенилфосфиннің анион-радикалы.....	102
<i>Баешова А.К., Молайган С., Баешов А.Б.</i> Суыткізгіш энергетиканың қазіргі замандағы жағдайы және суыткізгіш әдістері	107
<i>Закарина Н.А., Дәлелханұлы О., Жумадуллаев Д.А., Ақурпекова А.К., Джумабаева Л.С.</i> Al, AlZr және Ti-мен пилларирленген Na- және Ca-формалы монтмориллонитке енгізілген Pt- және Pd-катализаторларындағы тікелей айдалған бензиннің жеңіл фракциясының изомеризациясы.....	117
<i>Нәсіров Р.Н.</i> ЭПР спектроскопия көмегімен каспий маңындағы мұнайлардағы ванадийді анықтау.....	125
<i>Байжуманова Т.С., Тунгатарова С.А., Xanthopoulou G., Жексенбаева З.Т., Кауменова Г.Н., Еркибаева М.К., Жумабек М., Касымхан К.</i> Метанның олефиндерге дейін каталитикалық конверсиясы.....	132
<i>Калимукашева А.Д., Калиманова Д.Ж., Иманкулова З.А.</i> Формативті бағалау-химия сабақтарында оқыту процесінің ажырамас бөлігі.....	139
<i>Масенова А.Т., Калыкбердиев М.К., Сасс А.С., Кензин Н.Р., Канатбаев Е.Т., Цыганков В.П.</i> Бензин фракцияларындағы хош иісті көмірсутектерді жоғары қысымда отырғызылғын катализаторларды қолдану арқылы суыткендіру.....	146

СОДЕРЖАНИЕ

<i>Кантуреева Г.О., Defrancesco E., Алибеков Р.С., Уразбаева К.А., Ефимова И.Е.</i> Новые тенденции в идентификации традиционной пищевой продукции Казахстана	6
<i>Туктин Б.Т., Тенизбаева А.С., Нургалиев Н.Н., Шаповалова Л.Б., Яскевич В.И.</i> Исследование гидроочистки и гидроизомеризации прямогонной бензиновой фракции на модифицированных Ni(Co)-Mo- Al ₂ O ₃ - катализаторах	13
<i>Ахметалимова А.М., Ивасенко С.А., Марченко А.Б., Ишмуратова М.Ю., Полезчак Э., Людвичук А., Лосева И.В.</i> Исследование химического состава <i>THYMUS EREMITA KLOK.</i> и <i>THYMUS RASITATUS KLOK.</i> Карагандинского региона	20
<i>Фазылов С.Д., Нуркенов О.А., Журинов М.Ж., Аринова А.Е., Туктаров А.Р., Исаева А.Ж., Шахова Б.К.</i> Катализируемое комплексами палладияциклоприсоединение гидразонов к фуллерену C ₆₀	26
<i>Опимах Е.В., Левданский А.Э., Голубев В.Г., Корганбаев Б.Н., Сарсенбекулы Д.</i> Перспективные направления снижения удельных энергозатрат при измельчении	32
<i>Кансамет М.Ж., Тажиббаева С.М., Уракаев Ф.Х., Уралбеков Б.М., Буркитбаев М.М., Бачилова Н.В.</i> Получение и стабилизация наносеры	41
<i>Байсанов С.О., Толоконникова В.В., Нарикбаева Г.И., Корсукова И.Я., Жучков В.И.</i> Термодинамическая оценка выплавки марганцевых и хромистых ферросплавов на основе анализа их диаграмм состояния	47
<i>Кулекеев Ж.А., Нуртаева Г.К., Мустафин Е.С., Айнабаев А.А., Мустафин Т.Е., Борсынбаев А.С., Жарикесов Г.А.</i> Возможности использования хердеров при ликвидации разливов нефти на море	58
<i>Туктин Б.Т., Нургалиев Н.Н., Тенизбаева А.С., Шаповалова Л.Б., Комашко Л.В.</i> Гидрооблагораживание различных бензиновых фракций на модифицированных алюмокобальтмолибденовых катализаторах	67
<i>Калдыбекова А.Ж., Амангазиева А.Т., Халменова З.Б., Умбетова А.К.</i> Разработка технологии комплексного выделения биологических активных веществ из растений рода <i>Naplophyllum</i> A. Juss	74
<i>Опимах Е.В., Левданский А.Э., Волненко А.А., Жумадуллаев Д.К.</i> Методы проведения флотационных процессов	82
<i>Чиркун Д. И., Левданский А. Э., Волненко А.А., Сарсенбекулы Д.</i> Исследование динамики частиц в ударно-центробежных мельницах	92
<i>Баймукашева Г.К., Калауова А.С., Куспанова Б.К., Насиров Р.Н.</i> Анион-радикал трифенил-фосфина	102
<i>Баешова А.К., Молайган С., Баешов А.Б.</i> Современное состояние водородной энергетики и способы получения водорода	107
<i>Закарина Н.А., Дәлелханұлы О., Жумадуллаев Д.А., Акурпекова А.К., Джумабаева Л.С.</i> Изомеризация легкой фракции прямогонного бензина на Pt- и Pd-катализаторах, нанесенных на пилларированный Al, AlZr и Ti монтмориллонит в Na- и Ca-формах	117
<i>Насиров Р.Н.</i> Определение ванадия в нефтях прикаспийского региона методом ЭПР-спектроскопии	125
<i>Байжуманова Т.С., Тунгатарова С.А., Xanthoroulou G., Жексенбаева З.Т., Кауменова Г.Н., Еркибаева М.К., Жумабек М., Касымхан К.</i> Каталитическая конверсия метана в олефины	132
<i>Калимукашева А.Д., Калиманова Д.Ж., Иманкулова З.А.</i> Формативное оценивание - неотъемлемая часть процесса обучения на уроках химии	139
<i>Масенова А.Т., Калыкбердиев М.К., Сасс А.С., Кензин Н.Р., Канатбаев Е.Т., Цыганков В.П.</i> Гидрирование ароматических углеводородов в бензиновых фракциях на нанесенных катализаторах под давлением	146

CONTENTS

<i>Kantureeva G.O., Defrancesco E., Alibekov R.S., Urazbayeva K.A., Efimova I.E.</i> New trends in the identification of the traditional food products of Kazakhstan	6
<i>Tuktin B.T., Tenizbaeva A.S., Nurgaliyev N.N., Shapovalova L.B., Yaskevich V.I.</i> Study of hydro purification and hydroisomerization straight-run gasoline fraction over modified Ni(Co)-Mo- Al ₂ O ₃ - catalysts	13
<i>Akhmetlimova A.M., Ivashenko S.A., Marchenko A.B., Ishmuratova M.Yu., Poleszak E., Ludwiczuk A., Loseva I.V.</i> The study of the chemical composition of <i>THYMUS EREMITA</i> KLOK. and <i>THYMUS RASITATUS</i> KLOK. from the Karaganda region	20
<i>Fazylov S.D., Nurkenov O.A., Zhurinov M.Zh., Arinova A.E., Tuktarov A.R., Issayeva A.Zh., Shaihova B.K.</i> Catalyzed by palladium complexes the cycloaddition of hydrazones to fullerene C ₆₀ (in English).....	26
<i>Apimakh Ye.V., Leudanski A.E., Golubev V.G., Korganbayev B.N., Sarsenbekuly D.</i> Promising directions of reducing specific energy costs in grinding (in English).....	32
<i>Kapsamet M.Zh., Tazhibayeva S.M., Urakaev F.Kh., Uralbekov B.M., Burkitbayev M.M., Bachilova N.V.</i> Obtaining and stabilization of nanosulfur	41
<i>Baisanov S.O., Tolokonnikova V.V., Narikbayeva G.I., Korsukova I.Ya., Zhuchkov V.I.</i> Thermodynamic assessment of smelting of manganese and chromium ferroalloys based on the analysis of their state diagrams	47
<i>Kulekeyev Zh.A., Nurtayeva G.K., Mustafin E.S., Ainabayev A.A., Mustafin T.E., Borsynbayev A.S., Zharikessov G.A.</i> Using herders for oil spill response in the sea	58
<i>Tuktin B.T., Nurgaliyev N.N., Tenizbaeva A.S., Shapovalova L.B., Komashko L.V.</i> Hydrotreating of various petrol fractions over modified alumocobaltmolybdenic catalysts	67
<i>Kaldybekova A.Zh., Amangazyeva A.T., Halmenova Z.B., Umbetova A.K.</i> Development of technology for the complex isolation of biological active substances from plants of the genus <i>Haplophyllum</i> A. Juss	74
<i>Apimakh Ye.V., Leudanski A.E., Volnenko A.A., Zhumadullaev D.K.</i> Methods of carrying out flotation processes	82
<i>Chyrkun D.I., Levdanskiy A.E., Volnenko A.A., Sarsenbekuly D.</i> Study of the particle dynamics in impact-centrifugal mills (in English).....	92
<i>Baymukasheva G.K., Kalauova A.S., Kuspanova B., Nasirov R.N.</i> Triphenylphosphine anion radical.....	102
<i>Bayeshova A.K., Molaigan S., Bayeshov A.B.</i> Hydrogen energetics current state and hydrogen production methods.....	107
<i>Zakarina N.A., Dolelkhanyly O., Jumadullaev D.A., Akurpekova A.K., Djumabaeva L.S.</i> Isomerization of light fraction of straight-run gasoline on Pt- and Pd-catalysts supported on pillared by Al, AlZr and Ti montmorillonite in Na- and Ca-forms.....	117
<i>Nasirov R.N.</i> Determination of vanadium in the precaspian region's oil by the EPR-spectroscopy method.....	125
<i>Baizhumanova T.S., Tungatarova S.A., Xanthopoulou G., Zheksenbaeva Z.T., Kaumenova G.N., Erkibaeva M.K., Zhumabek M., Kassymkan K.</i> Catalytic conversion of methane into olefins.....	132
<i>Kalimukasheva A.D., Kalimanova D.Z., Imankulova Z.A.</i> Formative evaluation is an uninterrupted part of the training process on lessons of chemistry.....	139
<i>Massenova A.T., Kalykberdiyev M.K., Sass A.S., Kenzin N.R., Kanatbayev E.T., Tsygankov V.P.</i> Hydrogenation of aromatic hydrocarbons in gasoline fractions over supported catalysts under pressure.....	146

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://www.chemistry-technology.kz/index.php/ru/>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Т. А. Апендиев, Аленов Д.С.*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 11.10.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,8 п.л. Тираж 300. Заказ 5.